

# Recent Surveys in the Forests of Ulu Segama Malua, Sabah, Malaysia, Show That Orang-utans (*P. p. morio*) Can Be Maintained in Slightly Logged Forests

Marc Ancrenaz , Laurentius Ambu, Indra Sunjoto, Eddie Ahmad, Kennesh Manokaran, Erik Meijaard, Isabelle Lackman

Published: July 9, 2010 • DOI: 10.1371/journal.pone.0011510

## Abstract

### Background

Today the majority of wild great ape populations are found outside of the network of protected areas in both Africa and Asia, therefore determining if these populations are able to survive in forests that are exploited for timber or other extractive uses and how this is managed, is paramount for their conservation.

### Methodology/Principal Findings

In 2007, the “Kinabatangan Orang-utan Conservation Project” (KOCP) conducted aerial and ground surveys of orang-utan (*Pongo pygmaeus morio*) nests in the commercial forest reserves of Ulu Segama Malua (USM) in eastern Sabah, Malaysian Borneo. Compared with previous estimates obtained in 2002, our recent data clearly shows that orang-utan populations can be maintained in forests that have been lightly and sustainably logged. However, forests that are heavily logged or subjected to fast, successive coupes that follow conventional extraction methods, exhibit a decline in orang-utan numbers which will eventually result in localized extinction (the rapid extraction of more than 100 m<sup>3</sup> ha<sup>-1</sup> of timber led to the crash of one of the surveyed sub-populations). Nest distribution in the forests of USM indicates that orang-utans leave areas undergoing active disturbance and take momentarily refuge in surrounding forests that are free of human activity, even if these forests are located above 500 m asl. Displaced individuals will then recolonize the old-logged areas after a period of time, depending on availability of food sources in the regenerating areas.

### Conclusion/Significance

These results indicate that diligent planning prior to timber extraction and the implementation of reduced-impact logging practices can potentially be compatible with great ape conservation.

**Citation:** Ancrenaz M, Ambu L, Sunjoto I, Ahmad E, Manokaran K, Meijaard E, et al. (2010) Recent Surveys in the Forests of Ulu Segama Malua, Sabah, Malaysia, Show That Orang-utans (*P. p. morio*) Can Be Maintained in Slightly Logged Forests. PLoS ONE 5(7): e11510. doi:10.1371/journal.pone.0011510

**Editor:** Jerome Chave, Centre National de la Recherche Scientifique, France

**Received:** April 9, 2010; **Accepted:** June 12, 2010; **Published:** July 9, 2010

**Copyright:** © 2010 Ancrenaz et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This specific study was supported by a grant from the Australian Regional natural Heritage program and from the US Fish and Wildlife Services Great Ape Conservation Fund. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

The natural habitat of the orang-utan, the tropical forests of Sumatra and Borneo, are declining at an alarming rate as a result of human activities, such as agriculture and timber extraction. In Borneo, approximately ten percent of the remaining forests are protected for conservation, but it is doubtful that this network of protected areas alone will ensure the long-term survival of the species that leave in these forests [1], [2]. Early studies have suggested that the orang-utan was dependant on primary forests for survival and that forest exploitation and degradation was resulting in the rapid decline of the species [3], [4], [5], [6], [7], [8]. However, it is increasingly recognized that great apes (including orang-utans) can survive in low-impact and sustainably logged forests [9], [10], [11], [12], [13], [14].

Considering that more than 75% of the wild orang-utan populations in Borneo are currently found in forests that are exploited for timber [11], [15], understanding how orang-utan populations react and adapt to logging is becoming one of the major priorities for conserving the species at the landscape scale. Nevertheless, there is still a general lack of knowledge and information regarding how orang-utans respond to different intensities of timber extraction.

In 2002, our surveys in Sabah established that the commercial forest reserves of the Ulu Segama-Malua-Kuamut-Kalabakan complex were home to approximately 4,500 individuals, making it the largest unfragmented population of wild orang-utans in Malaysia [11]. These mixed lowland dipterocarp forests are located in the central part of the State and have been exploited for timber since the late 1950s [16]. Acknowledging the importance of the forests of Ulu Segama Malua (USM) for orang-utan conservation, the Sabah State government banned logging for a ten year period at the end of 2007. In 2006–2007, we conducted new aerial and ground surveys in these forests in order to monitor population trends; identify the primary proximate and ultimate factors impacting orang-utan abundance in disturbed forests; document fluctuations of orang-utan abundance in locations exposed to different logging intensities; and to propose that orang-utan conservation should be included in the forest management plan developed for this area.

In this paper we present data on the abundance and distribution of orang-utans in the forest reserves of the Ulu Segama-Malua region that were obtained during our 2006–07 surveys. We seek to determine whether the North Bornean orang-utan subspecies (*Pongo pygmaeus morio*) can maintain viable populations in sustainably and minimally logged forests, at least in the short-term? If so, does a threshold of habitat disruption and degradation exist, where maintaining a viable population becomes impossible? We discuss some of the possible reasons behind orang-utan resilience and we also provide general recommendations for maintaining populations in forests that are exploited for timber.

## Results

### Aerial transects

In May 2007, sixteen parallel lines interspaced by about 5 km were flown over Ulu Segama Forest Reserve for a total length of 344.4 km, and eight transects interspaced by approximately 2.5 km were flown over Malua Forest Reserve totaling 140 km (Map 1). Therefore, the total survey effort of aerial surveys was roughly 6% of the entire USM area. Land below 450 m asl accounted for 78.7% and land above 600 m asl for 5.8% of our sampling. Fair forest only represented 11.5%, and it was predominantly found on steep and higher ground located in the southern and western side of Ulu Segama, bordering the DVCA, and on the top of steep hills that were not accessible to heavy logging machinery. Highly degraded forests accounted for 52% of our sample and were characterized by the complete disruption of the original canopy structure; an extreme rarity of emergent climax trees; open areas; abundance of old/recent logging roads; and the presence of invading bushes, creepers and pioneer trees belonging to Euphorbiaceae and Rubiaceae families. We applied a correction factor of  $F_1 = 0.54$  to aerial indexes obtained in this overdegraded habitat to take into account the increased detectability and artificially inflated aerial indexes in overdegraded forests due to canopy openness [11]. Degraded forests accounted for 11.5%, and areas of active logging for 4.4% (mostly North Ulu Segama and Malua FR). Overall, a total number of 3199 orang-utan nests were recorded. In order to investigate if orang-utan abundance depended on topographical and connectivity features, we pooled the transects in several “Sampling Units”: North Ulu Segama (NUS); Segama East; Segama Central; Segama South and West; Malua. Within each sampling unit we then investigated possible fluctuations of density resulting from habitat differences.

**North Ulu Segama (NUS):** the forests of NUS cover roughly 12,000 ha. These forests are surrounded by oil palm plantations in the north and the Segama River in the south. They are highly degraded as a result of over-logging and fires. The last round of timber extraction was taking place in 2007 at the time of our surveys. In most areas the canopy was completely disrupted, few trees were left standing, logging roads and open areas were common, and pioneer trees such as *Macaranga spp.* dominated the landscape. Signs of active or very recent logging activities were widespread and distributed throughout the entire region. Approximately 40% of the aerial transects were flown over lands devoid of trees, considered as unsuitable habitat for orang-utans. Although orang-utan nests were found throughout the entire NUS area, they tended to be concentrated in compartments with the best forest stands or in isolated patches of trees found in the middle of over-degraded and open areas. Nests were slightly more abundant in areas with no active logging (Table 1).

Area	Transect	Length	Aerial Index	Habitat Type	Length (km)	Aerial Index	OU density (individuals/km <sup>2</sup> )	
North Ulu Segama	TUVWXY north	20.5	4.169	Overdegraded	11.4	4.77	1.98	
				Active logging	9.0	2.51	1.08	
	Segama East	ZYX	79.0	0.382	Overdegraded	42.8	8.291	0.12
					Degraded	36.2	8.412	0.18
Z		21.9	0.289	Active logging	2.1	8.218	0.11	
				Below 450 m asl	19.2	8.302	0.12	
				450-600 m asl	2.0	8.448	0.18	
Segama Central	TUVAW	147.6	1.068	Overdegraded	49.5	2.14	0.95	
				Degraded	47.1	2.44	0.97	
	U	41.0	2.729	Active logging	10.1	1.85	0.43	
				Below 450 m asl	19.7	1.41	0.57	
	T	65.6	2.032	Below 450 m asl	30.2	1.26	0.71	
				450-600 m asl	12.8	1.87	0.75	
				600 m asl	21.6	1.26	0.62	
Segama South West	WVUTPQR	92.7	6.07	Overdegraded	14.9	3.425	1.35	
				Degraded	7.0	6.145	2.48	
	Q	9.3	7.156	Macaranga	6.45	1.918	0.78	
				Active logging	2.8	2.424	0.97	
	D	29.0	3.031	Fair forest	21.5	5.656	2.23	
				Below 450 m asl	14.3	4.412	1.73	
W	18.1	6.022	DVCA	11.25	3.851	1.57		
			Below 450 m asl	6.05	3.299	1.27		
				450 m asl	11.8	4.812	1.83	
Malua	RUCDEFGH	140.0	4.168	Overdegraded	11.2	3.913	1.68	
				Degraded	122.5	4.408	1.76	
	F	18.0	3.288	Fair forest	3.5	6.118	2.41	
				Below 450 m asl	15.8	4.219	1.68	
	D	27.0	7.262	450 m asl	4.0	1.723	0.69	
C	36.0	5.234						
E	24.0	3.353						
K	9.0	7.289						

**Table 1. General results of aerial surveys in the five sampling units distinguished in the Ulu Segama Malua Forest Reserves.**  
doi:10.1371/journal.pone.0011510.t001

**Segama East/Central:** in eastern and central Segama, density increased from the eastern lines (ZYX pooled together: AI = 0.3 nest km<sup>-1</sup>) to the western lines (WVUT: AI = 2.0 nest km<sup>-1</sup>): t-test; t = 3.4; df = 5; p = 0.018\*. The forests of eastern Segama were highly degraded and harbored very low nest densities. Nests were more abundant in steeper, higher terrains that had patches of healthier forest and were located further away from active logging activities. In the central parts of Segama, orang-utan distribution was relatively uniform. Their abundance was lower in areas with active, on-going logging activities and maximum in regenerating and healthier forests located upland (>600 m asl), where logging activities occurred over ten years ago. However densities dropped

trastically in upland forests that recently experienced intensive logging (line T: 0.47 ind.km<sup>-2</sup>). In the past, tall trees were used by orang-utans to cross the Bole and Kawag Rivers, but these water bodies cannot be crossed by the animals following the removal of these trees. By considering the transect lines of the same habitat located on both sides of these rivers, we investigated local differences in nest distribution and abundance. Orang-utans were more abundant on the western rather than on the eastern side of the Bole River (West: L = 22.4 km of line, 139 nests, AI = 3.10 nest km<sup>-1</sup>, D<sub>OU</sub> = 1.23 ind. km<sup>-2</sup>; East: L = 23.2 km, 89 nests, AI = 1.91 nest km<sup>-1</sup>, D<sub>OU</sub> = 0.76 ind. km<sup>-2</sup>), and no difference was found on either side of the Kawag river, but orang-utan density was lower within the Kawag loop, which is more difficult for the animals to access: L = 17.9 km, 62 nests, AI = 1.73 nest km<sup>-1</sup>, D<sub>OU</sub> = 0.69 ind.km<sup>-2</sup>).

**Segama South/West:** Lines M and S border the south of the protected forests of DVCA in the westmost part of Segama. Part of length O was flown over DVCA and although data is presented in Table 1, it has not been included in our final analysis. Fewer orang-utan nests were identified in forests below 450 m asl (AI = 2.95 nest km<sup>-1</sup>) than above 450 m asl (AI = 5.23 nest km<sup>-1</sup>), with a difference that is near significance: U Mann and Whitney: z = -1.938: p = 0.053. The lowest nest densities in Segama S/W were recorded in lowland areas that have been highly disturbed by active and recent logging operations, and in areas highly invaded by *Macaranga sp.* Primary and old regenerating forests found in the highlands were the least disturbed habitat due to the steep slopes that are characteristic of this habitat, which limit and prevent conventional logging practices. These forests were home to the highest orang-utan densities recorded in Segama FR with about 2.1 ind. km<sup>-2</sup>.

**Malua FR:** Heavy logging occurred in Malua until the end of 2007, and most forests appeared degraded (43.4% of the total aerial length) or overdegraded (56.6%). Orang-utan abundance was higher in degraded (D<sub>OU</sub> = 1.76 ind. km<sup>-2</sup>) rather than in overdegraded forests (D<sub>OU</sub> = 1.00 ind. km<sup>-2</sup>). The highest density with about 2.4 ind. km<sup>-2</sup> was recorded in the forests of the “Bornean Biodiversity Conservation Plot” that appeared to be in very good condition. These forests are mature secondary forests and show a relatively diverse tree composition and structure. Orang-utan nests were more abundant on the western side of the Malua River (lines ABCD, 33.8 km, AI = 5.65 nests km<sup>-1</sup>, D<sub>OU</sub> = 2.21 ind. km<sup>-2</sup> versus lines CDEF, 57.9 km, AI = 3.97 nest/km, D<sub>OU</sub> = 1.57 ind. km<sup>-2</sup>) and in the northern region (D<sub>OU</sub> = 2.02 ind. km<sup>-2</sup>), than in the southern area (D<sub>OU</sub> = 1.20 ind. km<sup>-2</sup>) or in the overdegraded forests of the “Wildlife Corridor” that is located in the south-eastern part of Malua (D<sub>OU</sub> = 1.00 ind. km<sup>-2</sup>).

#### Ground transects and nesting sites

Line transects were conducted for ground truthing of the aerial data and for investigating local variations of orang-utan abundance. We performed a total of 106 ground line transects (total length of 79.4 km; average length: 742 m; range: 170–1710 m) over nine expeditions throughout USM between August 2006 and June 2007 (survey effort of 0.06%): Table 2. During this time we recorded a total of 1111 orang-utan nests built in a minimum of 35 families and 65 taxa of trees (18.0% of nesting trees were not identified, adding an unknown number of families and taxa as possible nesting site): Table 3. Four tree families and 4 genera represented 62.2% and 55.2% of all nesting sites respectively. When we considered the eight families most often used for nesting, we found no significant difference between choice of tree species for nesting and family abundance in the forest (values of Wilcoxon tests are given in Table 3). However, in highly disturbed forests, orang-utans preferentially used pioneer trees like *Neolamarckia cadamba* (40.1% in Malua and 19.4% in NUS) or *Pterospermum sp.* (4.0% and 38.4%). , In the contrary *Shorea sp.* (18.2%) and mature *Macaranga sp.* (13.7%) were preferentially used as nesting sites in the less degraded habitats or in the old-logged forests of Segama. For each survey site, we classified the forest into two major classes of habitat disturbance: degraded and overdegraded. Compared to degraded forests, overdegraded forests were characterized by: more logging roads (5.5 vs 3.4 roads/km of transect, although the difference was not significant: t-test value = 1.19; df = 8; p = 0.26); a significant lower basal area (8.0 vs 16.3 m<sup>2</sup>/ha: t-test = -5.51; df = 80; p<0.0001\*); and a significant lower tree density (142.7 vs 214.6 trees dbh>10 cm/ha: t-test = -3.85; df = 80; p = 0.0002\*). Variations of orang-utan density between overdegraded and degraded habitat were tested for in each expedition where the two types of habitat were present (n = 5). Densities were significantly higher in degraded (general average of 2.23 ind. km<sup>-2</sup>), versus overdegraded forests (1.36 ind. km<sup>-2</sup>): paired-sample t-test, df = 3, t = 6.79, p = 0.007\*. Ground truthing of aerial data was achieved by (1) pooling together all ground transects from different expeditions that were run in the same area, in similar habitat type where no significant difference in encounter rates were detected; and (2) comparing these with aerial orang-utan densities obtained over corresponding areas (n = 6 sites). Results given in Table 4 show a very strong correlation (r = 0.975) and no significant difference between the two data sets (paired-sample t-test: n = 6; t = 1.929; df = 5; p = 0.11).

XPDC	Location	Type	LTs	Length (km)	Altitude (m)	Basal Index	Basal Area (m <sup>2</sup> /ha)	Tree Density (per ha)	Orang-utan density (CI)
1	Malua South	Deg.	5	4.515	250-400	9/9	9/9	878	1.35 (0.40)
2	Malua NW	Overdeg.	2	4.284	250-400	9/9	9/9	878	0.69 (0.23)
		Overdeg.	2	2.180	200-400	9/2	4/1 (2)	90	2.19 (0.7)
		Overdeg.	2	1.080	200-400	2/2	2/1 (2)	90	1.89 (0.6)
3	Malua NE	Overdeg.	1	4.841	200-400	1/1	1/1 (0)	212	1.62 (0.5)
4	Segama SW	Overdeg.	3	4.889	200-400	7/1	11/1 (0)	1123	1.21 (0.4)
4-5	Segama SW	Overdeg.	8	4.841	200-400	2/2	21/2 (7)	294	1.8 (0.5)
6	Segama SW	Overdeg.	13	4.876	200-400	5/5	10/2 (7)	315	0.4 (0.4)
6	Segama SW	Overdeg.	8	5.011	200-400	5/1	11/2 (0)	301	2.1 (0.2)
7	Malua NW	Overdeg.	7	6.813	200-400	9/4	9/8 (0)	307	0.7 (0.2)
8	Malua NE	Overdeg.	9	6.516	200-400	3/5	9/8 (0)	121	1.61 (0.4)
9	Malua NE	Overdeg.	14	11.510	200-400	9/2	8/2 (1)	305	2.22 (0.2)
9	NUS	Overdeg.	13	6.311	200-400	2/1	4/1 (1)	189	2.32 (0.4)

Legend: Deg = degraded; Overdeg = overdegraded; alt = above sea level; NW = north-west; NE = north-east; SW = south-west; CI = coefficient of variation obtained by DeSmet.

doi:10.1371/journal.pone.0011510.t002

**Table 2. Location, main characteristics, orang-utan densities (with associated Coefficient of Variation) of all ground surveys conducted in the USM forests.**

doi:10.1371/journal.pone.0011510.t002

Number	Total		Malua		Segama		NUS	
	Nests	Trees	Nests	Trees	Nests	Trees	Nests	Trees
<b>Dipterocarpaceae</b>	<b>11.1</b>	<b>28.0</b>	<b>12.6</b>	<b>26.2</b>	<b>22.4</b>	<b>26.9</b>	<b>6.1</b>	<b>11.3</b>
Malua sp.	11.1	28.0	8.6	18.5	16.7	24.8	6.7	7.7
Other spp.	0	0	4.0	7.7	5.7	2.1	0	3.6
<b>Euphorbiaceae (Dipterocarp sp.)</b>	<b>1.3</b>	<b>3.3</b>	<b>1.3</b>	<b>3.1</b>	<b>2.0</b>	<b>2.3</b>	<b>0.0</b>	<b>3.5</b>
Alseodaphne sp.	1.3	3.3	1.3	3.1	2.0	2.3	0.0	3.5
Other spp.	0	0	0	0	0	0	0	0
<b>Euphorbiaceae</b>	<b>19.8</b>	<b>49.4</b>	<b>6.7</b>	<b>16.1</b>	<b>15.2</b>	<b>9.8</b>	<b>6.6</b>	<b>16.1</b>
Alseodaphne sp.	6.7	16.1	6.7	16.1	15.2	9.8	6.6	16.1
Other spp.	13.1	33.3	0	0	0	0	0	0
<b>Fagaceae (Lathyrus sp.)</b>	<b>4.4</b>	<b>1.8</b>	<b>5.3</b>	<b>1.7</b>	<b>16.8</b>	<b>2.8</b>	<b>1.0</b>	<b>0</b>
Lathyrus sp.	4.4	1.8	5.3	1.7	16.8	2.8	1.0	0
Other spp.	0	0	0	0	0	0	0	0
<b>Myrtaceae (Palaquium sp.)</b>	<b>1.1</b>	<b>0.8</b>	<b>1.0</b>	<b>0.6</b>	<b>1.2</b>	<b>0.8</b>	<b>1.4</b>	<b>0.7</b>
Palaquium sp.	1.1	0.8	1.0	0.6	1.2	0.8	1.4	0.7
Other spp.	0	0	0	0	0	0	0	0
<b>Myrtaceae (Lagotis sp.)</b>	<b>2.0</b>	<b>3.0</b>	<b>5.7</b>	<b>8.1</b>	<b>6.8</b>	<b>8.4</b>	<b>2.8</b>	<b>5.8</b>
Lagotis sp.	2.0	3.0	5.7	8.1	6.8	8.4	2.8	5.8
Other spp.	0	0	0	0	0	0	0	0
<b>Palisotaceae</b>	<b>24.5</b>	<b>8.8</b>	<b>42</b>	<b>16.1</b>	<b>3.3</b>	<b>8.8</b>	<b>21.8</b>	<b>15.3</b>
Palisotaceae sp.	24.5	8.8	42	16.1	3.3	8.8	21.8	15.3
Other spp.	0	0	0	0	0	0	0	0
<b>Sterculiaceae (Pterocarpus sp.)</b>	<b>12.6</b>	<b>7.3</b>	<b>8.7</b>	<b>3.6</b>	<b>4.0</b>	<b>3.7</b>	<b>16.4</b>	<b>29.4</b>
Pterocarpus sp.	12.6	7.3	8.7	3.6	4.0	3.7	16.4	29.4
Other spp.	0	0	0	0	0	0	0	0
<b>Other tree families</b>	<b>28.5</b>	<b>49.9</b>	<b>34.7</b>	<b>39.6</b>	<b>37.5</b>	<b>47.7</b>	<b>18.4</b>	<b>24.5</b>
Other tree families	28.5	49.9	34.7	39.6	37.5	47.7	18.4	24.5
Nikolov test values	Z = -8.054, p = 0.001		Z = -6.652, p = 0.015		Z = -1.601, p = 0.108		Z = -0.829, p = 0.403	

**Table 3. Percentage of utilization of the eight most common tree families and taxa used for nesting and percentage of tree abundance recorded in 69 botanical plots in three different areas: Malua, Segama and North Ulu Segama.**  
doi:10.1371/journal.pone.0011510.t003

Areas	Estimated ground density	Estimated aerial density
Kawag (4–5 <sup>th</sup> XPDC)	1.1	0.8
Silviculture area (3 <sup>rd</sup> XPDC)	1.2	0.8
West Segama (6 <sup>th</sup> XPDC)	2.7	2.4
South Malua (1 <sup>st</sup> ; 7 <sup>th</sup> XPDC)	1.1	1.2
North Malua (2 <sup>nd</sup> ; 6 <sup>th</sup> SPDC)	2.05	2.0
Sebagaya (4 <sup>th</sup> XPDC)	0.1	0.1

**Table 4. Orang-utan density estimates achieved during ground and aerial surveys over corresponding areas.**  
doi:10.1371/journal.pone.0011510.t004

The total orang-utan population size living in USM was obtained by combining the knowledge gained from aerial and ground transects and by following the stratification pre-established from our aerial lines: Table 5, Fig. 2. Our final estimate is that there are 2,580 orang-utans (968–7275) in the forests of Ulu Segama Malua.

Area Code	Size (km <sup>2</sup> )	Location	Density	95% CI	Orang-utan Number	95% CI
1	16.24	Sebagaya	0.00	0.0–0.015	1	0–2
2	23.04	WGA	0.00	0.0–0.015	2	0–5
3	891.26	East Ulu Segama NW 7/02 – Sabah – west RW 1/02 RW 3/01	0.15	0.0–0.4	12	10–172
4	51.81	Central RW 7/02	0.1	0.1–1.12	21	7–96
5	88.08	North Kuning Region – Kuning Area – SW 3/01 and RW 7/01	0.1	0.2–1.82	68	10–148
6	216.78	East Ribu Area	0.8	0.2–2.19	173	60–474
7	402.88	West Ribu Area – West Ribu Corridor – South Malua	1.1	0.4–5.8	136	107–1382
8	193.72	South Ribu Area – West RW 3/01 – East RW 7/02	1.2	0.44–5.27	181	68–491
9	102.28	South Ulu Segama – RW 7/02 – DCVM buffer	0.5	0.11–2.46	158	62–448
10	115.1	North Ulu Segama – South RW 3/04	1.5	0.55–4.81	172	64–422
11	343	South West Ulu Segama – RW 7/09 – DCVM buffer	1	0.71–5.47	158	146–1364
12	175.2	North Malua	2	0.71–5.47	346	121–940
13	50.48	West Malua – RW 3/03	1.0	0.56–4.36	91	29–228
14	56.08	South-west Malua	2	0.71–5.46	112	41–236
15	2.12	Sabah Biodiversity Park	2.4	0.81–6.59	7	2–16
<b>TOTAL</b>					<b>2289</b>	<b>1289–9686</b>

**Table 5. Number of orang-utans living in the USM forests estimated from the combination of ground and aerial surveys (See Figure 2 for the exact locations of the areas).**  
doi:10.1371/journal.pone.0011510.t005

## Discussion

It is now well established that estimating great ape abundance from nest densities can yield highly imprecise results due to the fluctuation in the nest decay rate values, amongst other factors [17]. Repeated nest counts are one way to reduce imprecision but the time required in these exercises further reduce the areas being investigated by surveyor teams [18]. In addition, monitoring the large areas that are typically occupied by great ape populations require major efforts that are difficult to match in the field due to financial, human and time constraints. Aerial surveys offer an interesting and cost-effective alternative to monitor orang-utan populations at the landscape level [11].

Combining ground and aerial surveys achieved a precise knowledge about the distribution, abundance and some of the factors influencing the orang-utan population living in the degraded forests of Ulu Segama Malua. Aerial surveys increased the general survey effort to 5.8% of the entire survey area, which is one of the highest scores documented for great apes [19] and is above the limit of 0.26%, recently proposed to achieve reliable nest abundance estimates [20]. A strong correlation was obtained between aerial and ground results, further validating the model recently developed in Sabah [11]. The discrepancy between aerial and ground indices identified in the forests of North Ulu Segama (NUS) was explained by the extreme degradation of this habitat. Because this area is a mosaic of trees left standing in bare land, ground line transects were predominately located in forested areas, while bare landscapes were typically avoided, in order to optimize time spent in the field. Therefore, orang-utan estimates were only available for forested areas and achieved a high score of 2.72 individuals/km<sup>2</sup>, without considering unsuitable habitat. On the contrary, aerial surveys covered all habitat types, forested or not, which resulted in an overall lower density (1.52 ind. km<sup>-2</sup>) compared to the ground data. Since our flights indicated that only 60% of the habitat was suitable for orang-utans, we used this stratification factor and ended up with similar population size estimates for NUS for ground (2.72×0.6×120 km<sup>2</sup> = 194 orang-utans) and aerial data (1.52×120 = 182 orang-utans).

Our surveys in 2007 in the forests of USM yielded similar population estimates (2,600 individuals) to our 2002 surveys (2,300 individuals; 95% confidence intervals between 1,744 and 3,657), indicating that this population has been relatively stable over this five-year period. However, this general picture hides fluctuations that are occurring at geographical and local scales throughout the

entire landscape. The uneven orang-utan abundance in USM results from the interaction of historical, man-made and natural features.

#### Orang-utan abundance and human history

The scarcity of orang-utan nests identified in the eastern forests of USM (densities comprised between 0 and 0.2 ind. km<sup>-2</sup>) can be related to the regional human history. Eastern Sabah has been inhabited for approximately 30,000 years, based on the earliest signs of human occupation in the State [21], and small human communities have been permanently established along the lower part of Segama River for centuries [22]. During this time, people were venturing into the upper parts of Segama for hunting expeditions and to take refuge in times of trouble and epidemics [23]. Trading with China and other nations blossomed in the 15<sup>th</sup> century, targeting forest products and animal parts (rhinoceros horns, nests of swiflets, hornbill skulls etc.). Hunting orang-utans for meat, traditional medicine and for skulls (after the ban of human head-hunting), might have taken its toll on the original population in this area, and possibly led to local extinction. Currently, orang-utan densities in lower Segama are at their lowest close to well-established villages [16], as has been shown for other orang-utan populations that are subjected to hunting pressure [24]. Orang-utans are slow breeders and any given population will go extinct if the yearly hunting level exceeds 1% of a particular population [25]. In addition to the probable impact of past hunting pressure, our botanical plots revealed that the eastern forests of USM were heavily disturbed, which was identified by the lack of medium and large sized trees, a low basal area, the over-abundance of pioneer tree species and the extreme rarity of sizeable dipterocarp trees and other mother trees. These findings indicate that past fires or clear-cutting during previous logging cycles have ravaged these forests and may have wiped out local orang-utan sub-populations.

#### Orang-utan abundance and natural features

Orang-utan densities were higher in the west than in the east, and reached 2.0 to 2.5 ind./km<sup>2</sup> in some parts of Malua and southwest Segama. However, very few nests were recorded in limestone habitats and in forests growing on ultra basic soils originating from Bidu Bidu and similar associations. The lack of sodium and the relative abundance of nickel, chromium and cobalt characteristic of these soils limits the growth of many plant species, resulting in a less diverse tree community with fewer food resources than other forest types, which accounts for the lower nest abundance in these suboptimal orang-utan habitats [26]. Large bodies of water, such as the Segama River, represent a barrier to orang-utans dispersal [27], [28]. Orang-utan abundance showed differences on both sides of the Malua, Bole and Kawag Rivers, indicating that these bodies of water may act as potential barriers for dispersal following felling of large trees that originally acted as natural bridges (Figure 2; Table 5). In Borneo, orang-utan densities usually decrease with altitude and drop sharply above 500 m asl [5], [29], [30], [31]. However in USM, high concentrations were locally recorded above this threshold in several areas (2.7 ind./km<sup>2</sup> in Segama Central; 1.8 ind./km<sup>2</sup> in Segama SW), while densities were significantly lower in surrounding lowland forests (Table 1). In most cases, logging activities had recently occurred or were taking place concurrently to our surveys in the surrounding lowland forests. We can therefore hypothesize that logging resulted in the forced migration into the less preferred highland forests, where orang-utan populations sought “refuge” by leaving disturbed areas [11], [29]. In large parts of southwest USM with no signs of recent logging in surrounding lowlands, valleys and ridges located above 450 m asl harbored about 2 individual km<sup>-2</sup>. These highland forests were lightly logged 15 years ago, but their tree diversity, size, height and canopy cover achieved the best scores among all of the survey sites, indicating that these are mature habitats or forests in a healthy regeneration stage. Oak trees belonging to the Fagaceae family (*Lithocarpus* spp.) were particularly common and were fruiting shortly before and during surveys. Acorns are one of the favored orang-utan foods (Russon et al, 2008), and their abundant production could have attracted animals from lower lands, possibly explaining the seasonally inflated orang-utan abundance in these hills (Singleton, 2000).

#### Orang-utan abundance and logging

Overall, the habitat found in USM is very heterogeneous. This is a result of conventional logging practices and heavy extraction rates, coupled with high road densities and indiscriminate felling, that has led to a mosaic of highly degraded forests bordering hilltops and isolated patches of moderate habitat in a chaotic pattern. This condition results in extremely patchy and uneven nest distribution resulting in the large variation in nest encounter rates between ground transects located in the same survey areas, or in aerial scores fluctuating by more than 30 nests between two successive observation periods of 30 seconds (representing a distance of about 500 m). Orang-utans feed on a wide range of plants [32], but their density is limited by the frequency and duration of periods of food shortage and is correlated with fruit abundance during periods of low fruit availability [33]. Therefore, sites experiencing extreme periods of food shortages support lower population densities. In response to fruit shortages orang-utan's shift their diet to non-fruit sources and more fibrous vegetation like leaves and barks [5], and alter their range patterns in order to exploit alternative food resources [34]. In Borneo, forests dominated with dipterocarps experience extreme temporal fluctuations in fruit availability; dipterocarp abundance is negatively correlated with orang-utan abundance at many sites [14]. Sustainable and selective logging typically targets a small number of trees that are primarily Dipterocarps. After felling, timber species are replaced by pioneer and asynchronous trees (such as *Dracontomelon* sp., *Ficus* sp., *Neolamarckia cadamba*, etc) and light-demanding woody climbers (*Spatholobus* sp., *Uncaria* sp., etc). By fruiting more frequently than climax tree species and by providing young leaves and bark, these pioneer plants are supplying new and alternative food sources that buffer periods of food scarcity. In addition, exploited habitats experience changes in fruiting event patterns and species such as *Garcinia* sp. and *Litsea* sp., which are part of the orang-utan diet, will bear more fruit during this time, providing additional resources to the animals [33].

Our results show that in USM, lightly logged forests supported relatively high orang-utan densities that were occasionally higher than those encountered in some primary lowland mature forests (see Table 1). Forests that were only logged once, over 15 years ago, supported the highest orang-utan densities during our surveys, showing that orang-utans recolonize old regenerating forests and can re-establish densities similar to or even exceeding pre-logging conditions [10], [14]. Densities documented close to the Bole River during our surveys (around 2 ind./km<sup>2</sup>) are comparable with orang-utan abundance documented when these forests were still pristine [29], indicating that orang-utans have maintained their numbers in this area even though it has been subjected to 40 years of logging activities. However, forest patches with active disturbance systematically yielded lower orang-utan densities than surrounding forest that were not exploited at the time of our surveys, suggesting that the animals take refuge in less disturbed areas as suggested by Mac Kinnon [29]. Recolonization of previously logged areas will depend on the intensity of logging activities and the regeneration dynamic of the forest. The two most abundant pioneer trees identified during our surveys were *Macaranga* sp. (Euphorbiaceae) in Segama and *Neolamarckia cadamba* (Rubiaceae) in Malua. *Macaranga* colonizes quickly in clear-cut areas and old logging roads and has the ability to outgrow other tree species, resulting in sizeable pure stands in the most degraded areas. These trees produce wind-dispersed seeds and offer very little food resources to the fruit eating community. *N. cadamba* on the contrary, produces both fruit and bark that are edible and consumed by orang-utans. In Malua, we recorded numerous signs of bark consumption and a huge proportion of nests built in these trees (Table 3). *N. cadamba* with its spaced crown also does not

restrain other trees from colonizing the areas, which helps to maintain a more bio-diverse forest within a localized area. Orang-utan density was significantly higher in areas of *N. cadamba* growth than in *Macaranga* dominated areas (1–2 individuals km<sup>-2</sup> versus 0.1–0.4 ind km<sup>-2</sup>). Because of lower food availability in *Macaranga* dominated regions, orang-utans have to forage over a much larger area, which results in lower densities in these forests.

At all survey sites, extremely damaged habitats yielded fewer nests than lightly logged forests (Table 1). Mechanical logging inflicts structural and incidental damages to all tree size-classes [35], and heavy logging results in impoverished forest composition (fewer tree diversity, fewer food sources) and structure (lower tree density, basal area and canopy height, absence of tall trees and contiguous canopy). The destruction of fallback food sources such as *Ficus sp.* and other key plant species in overlogged areas further impoverishes the habitat and induces significant orang-utan population decline. Simplification of the forest and destruction of the original forest mosaic, force orang-utans to either use a larger range or to adopt a “sit and wait” strategy to save energy and to digest more fibrous food [36]. When food resources are destroyed over large areas, this leads to a drastic decline, as documented for the orang-utan sub-population found in the NUS area. This sub-population is completely isolated from the main population by large oil palm plantations and by the Segama River. In 2002 before the latest logging cycle, the NUS forests were already highly degraded as a result of past fires and logging activities but they were still home to approximately 400 individuals [11]. Whereas in 2007, our estimates found that there were less than 200 animals in these same forests. This decline was due to the most recent logging cycle, which left an extremely degraded habitat with acutely low tree densities and basal areas, extensive openings in the canopy and very few food resources besides the leaves and bark of pioneer plants.

Eastern Borneo suffers the most from the El Nino Southern Oscillation events and from the resulting droughts, fires and periods of food scarcity [37], [38]. As a result, orang-utans have to survive on alternative food sources such as barks and leaves for extensive periods of time. Some anatomical features of the North eastern Bornean subspecies *P.p.morio* (more robust jaw bones, thicker teeth enamel, smaller skull size) could possibly be related to the specific ecological traits of the region [39]. It has been hypothesized that these anatomical features may predispose this subspecies to cope better with habitats with fewer fruit sources that have abundant fibrous fallback foods that are typical of lightly exploited forests [14], [40].

However, the wide dietary flexibility of the species as shown by the number of plants included in their diet [32], their intimate knowledge of the forest and their faculty of storing fat when food is abundant in the forest [41] are major factors accounting for the possible adaptation of the species to some level of habitat disturbance.

#### Conclusion

Our surveys in USM show that orang-utan populations can be maintained in lightly and sustainably logged forests but decline and are eventually driven to localized extinction in forests that are heavily logged or subjected to fast, successive coupes following conventional extraction methods. For example, the rapid extraction of more than 100 m<sup>3</sup> ha<sup>-1</sup> of timber, led to the crash of the sub-population found in the NUS forests.

Considering that the majority of great apes are currently found outside of protected areas in Africa and in Asia, it is clear that conserving these iconic species requires the establishment of a viable network of protected forests among a mosaic of certified logging concessions and other suitable habitats [13], [42]. However the orang-utan conservation community is not necessarily ready to seriously support the idea of conserving orang-utans in working timber concessions. This is partly due to the strong belief, to some extent based on empirical evidence, that all logging harms orang-utans [4], [14], and that therefore conservation of the species in timber concessions was not an option. Another factor may be that many conservationists find it difficult to entertain the notion of protecting a species in a forest that is not managed primarily for conservation purposes [43], even if orang-utans can survive in such habitats. We emphasize that with the majority of orang-utans occurring outside protected areas, and often in timber concessions, long-term protection of the species will require working closely with the timber industry and with concession managers. Not only are such programs more cost-effective than establishing new protected forests, they are also more likely to get support from government and other stakeholders [44].


To make orang-utan conservation in timber concession work, several crucial steps are required. The key recommendations for reconciling logging practices and orang-utan conservation, is the creation, preparation and implementation of a precise, pre-harvesting conservation management plan, in order to reduce incidental damages during felling and the adoption of reduced-impact logging practices (following the “Forest Stewardship Council” or another internationally recognized body) that will be followed during all exploitation phases. Priority will be placed on a strictly enforced, zero-hunting policy, as it has been shown that the poaching of orang-utans (or other great apes) will inevitably lead to population extinction. Crucial ecological resources (like riparian forests or patches of rich lowland forest ) need to be identified and set aside from exploitation compartments, and major food resources (like large fruit tree and key staple food plants like *Ficus sp.* and large climbers) must be protected from possible damages. Compartments to be exploited should follow a rotation system that allows for the creation of “refuge areas” that can be used by animals when they leave the vicinity experiencing active disturbance. Areas badly impacted by extraction activities (like stumping grounds or major logging roads) need to be rehabilitated and replanted with a mixed array of fast-growing fruit and timber trees, in order to enhance food resources in the forest. In silviculture treatment areas, large woody climbers that produce leaves and bark that are consumed by orang-utans and enable them to move throughout the forest, should not be cut. Finally, a thorough orang-utan bio-monitoring program must be developed and implemented to document population trends and their fluctuations in response to different logging regimes. If such management practices become standard practice in all remaining unprotected orang-utan habitats, chances of long-term survival of orang-utans in the wild would significantly increase.

#### Materials and Methods


##### Physical features of Ulu Segama Malua

The block of Ulu Segama/Malua (USM) is located in south central Sabah between 116°28'E and 4°14'N: it comprises the commercial forest reserves of Ulu Segama (202,856 ha) and Malua (33,969 ha), as well as four protected virgin jungle forests totaling 4,273 ha (Figure 1). The USM landscape is primarily below 600m asl and consists of a variably steep terrain, with many hills and ridges that reside over the plains which are typically located close to the large rivers (Segama, Bole, Kawag). Slopes above 25% cover 10% of the entire USM system and steep hills reaching 1250 m asl are concentrated in the areas surrounding “Danum Valley Conservation Area” (DVCA). Malua's topography is predominately flat and low except for a lone hill reaching roughly 700 m asl on its western flank and the higher land located close to DVCA. Geologic formations in USM include: crystalline basement originating from metamorphic and igneous rocks of the lower Triassic which are predominately located along rivers; Chert-Spilite sandstone from the late Cretaceous; and younger sedimentary and volcanic rocks from the formation of the Kuamut

area. Most of the soils are acidic, with a pH ranging from 3.6 to 5.4, are easily eroded and lose nutrients rapidly when they are disturbed. The wet tropical climate is created by the Indo-Australian monsoon system, the average annual temperature is 27°C (with an average maximum and minimum temperature of 31°C and 23°C respectively), and rain is brought in by the northerly winds from December to March. From 1976 to 1996, annual rainfall varied between 1775 mm to 3708 mm, with regular deficits every few years (El Nino). These droughts can be severe and result in intense tree mortality and fires [45].



**Figure 1. Location of aerial and ground surveys in Ulu Segama Malua, Sabah, Malaysian Borneo.**  
doi:10.1371/journal.pone.0011510.g001



**Figure 2. Orang-utan densities in the different areas of Ulu Segama Malua forests.**  
doi:10.1371/journal.pone.0011510.g002

#### Forest exploitation and forest types

Three major forest types occur naturally in USM and all are dominated by dipterocarp trees. The Lowland Mixed Dipterocarp Forest (LMDF) is typically found below 500 m asl. Common families include Dipterocarps (representing at least 60% of the basal area and 30% of the total tree density), Tiliaceae and Sapotaceae. Upland Mixed Dipterocarp Forests (above 500 m) are characterized by the abundance of Fagaceae (oak trees). Ultramafic Forests are found on the ultra-basic soils of the Bidu-Bidu formation that are deficient in phosphorus, potassium and calcium, are floristically less diverse and are of lower stature than other forest types. Throughout the entire landscape, the original forest structure and integrity has been altered drastically following multiple logging cycles that began in 1957 and finished at the end of 2007. The first round of logging (1957–1999), produced roughly 22 million m<sup>3</sup> logs from Ulu Segama (estimated production of 87.5 m<sup>3</sup>/ha) and 2.5 million m<sup>3</sup> logs from Malua (65.5m<sup>3</sup>/ha). The second round of logging, which was initiated in 1999 and completed in 2007, concentrated on approximately 105,000 ha in Ulu Segama and 20,000 ha in Malua and produced a significantly lower yield of 46.5 m<sup>3</sup>/ha in Ulu Segama and 33m<sup>3</sup>/ha in Malua [16]. Only a few protected areas escaped logging activities, namely DVCA and parts of Kawag Gibong and Sepagaya Virgin Jungle Forest Reserves. In addition, some of the logged forests were damaged by fires during major droughts induced by El Nino events. More than 5,000 ha were burned in Northern Ulu Segama (NUS) in 1983. In general, burnt areas become open land that are devoid of trees and are dominated by herbs and shrubs, with few or no signs of natural forest regeneration. However, patches of isolated, fire resistant, and regenerating pioneer species and trees from the old forest can be interspaced in this landscape [21].

The USM forests are currently classified into five different classes based on the density of trees >40 cm dbh (diameter at breast height), estimated from their crown size and visual interpretation of color aerial photographs. This approach provides a quick assessment of the potential commercial timber value of a forest [16]. The central part of USM forests (198,000 ha or 82%) are classified as very poor (i.e. less than 10 trees >40 cm dbh/ha, yielding less than 20 m<sup>3</sup>timber/ha), while about 23,000 (12%) ha are considered moderate to poor strata (between 10 and 30 trees 40 cm dbh/ha) and only 5% (or 13,000 ha) are considered good forest. Overall, more than 3% of USM is completely open and devoid of trees, and canopy cover is less than 30% in over 70% of USM. Low-lying and easily accessible areas have been particularly damaged by extensive and repeated coupes. These areas are characterized by: prolific signs of past human exploitation (tractor roads, stumping grounds, erosion, etc); an extremely low basal area and low tree density; a highly disrupted canopy with large forest gaps; and an abundance of invasive and pioneer plant species.

#### Survey methodology

The USM surveys combined ground and aerial data collection. Ground transects were randomly located on topographical maps (1:50 000) then ran on the ground. When possible, their direction was roughly perpendicular to large rivers, hills and major roads, in order to reduce between-transect variations and to achieve more reliable density estimates [46], [47]. Transect length was directly determined using a walking-distance measuring device: along each transect a team of two cleared a straight line-path and

confirmed the bearings with a compass, while a second team of three recorded information on forest type, general level of habitat degradation and nest presence. For each nest observed, we measured the perpendicular distance from the transect and recorded size, dbh, species of the nesting tree, as well as its approximate age [19]. Botanical plots (10×50 m) were randomly selected along the transects in order to characterize forest structure and composition. In all of the plots we identified the family or taxa levels of the trees with a dbh>10cm, recorded canopy height, climbers abundance, forest type and disturbance level (degraded and overdegraded).

Aerial transects followed randomly stratified parallel lines. We used a small helicopter, type Bell 206 Jet Ranger, and followed the methodology developed in Sabah and described by [11]. Helicopter speed and height were constant at 70 km/hour and 60–80 meters above the forest canopy. The co-pilot recorded the exact location of the aircraft every 30 seconds with a GPS, and collected information on habitat types, signs of wildlife presence and human activities. Combining available historic logging information and direct observation from the aircraft, we distinguished six different habitat classes: 1. Active logging; 2. Highly degraded and recently logged forests (logging less than 2 years old, many logging roads, open canopy, no re-growth of pioneer species); 3. Degraded forests (logging activities more than two years old, signs of forest regeneration, open canopy); 4. Old logged forests (logging more than 10 years old, most places –especially old logging roads- are infested with *Macaranga spp.* or Rubiaceae trees, canopy typically closed but short); 5. Fair forest (primary forest or forest lightly logged long ago, closed and diverse canopy, presence of diverse and tall emergent trees); 6. Unsuitable orang-utan habitat (river, large open areas, etc). From the back seat, two observers looked for orang-utan nests from either side of the helicopter and relayed all sightings to a nest recorder seated between them. All visible nests were recorded in this manner and the nest recorder noted the number of nests detected by the observers per each 30-s period.

#### Calculations

Ground nest densities were analyzed following line-transect analysis guidelines and were computed using the software Distance 4.2 [48]. Transformation of nest density into an orang-utan density was achieved with the formula:

$$\hat{D}_{Ape} = \frac{\hat{D}_{nest}}{\hat{p} * \hat{r} * \hat{t}}$$

where  $D_{nest}$  is the estimated nest density,  $p$  the estimated proportion of nest builders,  $t$  the estimated nest decay rate, and  $r$  the estimated daily rate of nest production. We used the same parameters previously determined in Sabah, to allow for direct comparison of population trends. We also used a differential nest decay rate taking into account the species of trees used for nesting [19].

Aerial indexes (number of nest per km of flight) were converted into nest and orang-utan density estimates with the model designed for Sabah and extensively described in Ancrenaz *et al.*, 2005:  $\hat{D}_0 = \exp[4.7297 + 0.9796 \log(AI_0)]$ ,  $\hat{D}_0$  being the estimated orang-utan nest density and  $AI_0$  the general aerial index ( $AI_0 = (AI_{left} + AI_{right})/2$ ). A final confidence interval for the predicted orang-utan

nest densities was achieved with  $\left[ \left[ \left( \frac{\hat{D}_0}{C}, \hat{D}_0 \times C \right) \right] \right]$ , where  $C = \exp(0.6067 \times \sqrt{1 + v^*})$  and  $v^* = 0.1908 - 0.2628 \times \log(AI_0) + 0.1132[\log(AI_0)]^2$ . Further statistical analysis were computed with the software SPSS.

#### Acknowledgments

For permission to work in Sabah we are grateful to the Economic Planning Unit, to late Datuk Patrick Andau (previous Director of the Sabah Wildlife Department) for his continuous support, to staff of the Sabah Forestry and Sabah Wildlife Department who were involved in these surveys. We would also like to thank Katie King and anonymous reviewers for their useful comments on the manuscript. Long-term KOCP activities have been supported by several zoos (Apenheul, Beauval, Cleveland, Columbus, Houston, La Palmyre, Oregon, Philadelphia, Phoenix, Saint Louis, Utah Ogle, Woodland Park); foundations (Abraham, Arcus, BOS-NL, Earth Shared, Elephant Family, Ensemble, Mohamed bin Zayed, Shining Hope, WoodTiger, World Women Work, Wildlife Conservation Network); US Fish and Wildlife Services, North England Zoological Society. Last but not least, the KOCP would not exist without the commitment of the 40 research assistants employed by the project who are dedicated to conserve the natural resources of the Kinabatangan floodplain and we are extremely grateful to all of them.

#### Author Contributions

Conceived and designed the experiments: MA LA KM. Performed the experiments: MA IS EA KM. Analyzed the data: MA EA KM. Wrote the paper: MA EM IL.

#### References

1. Curran LM, Trigg SN, McDonald AK, Astiani D, Hardiono YM, et al. (2004) Lowland forest loss in protected areas of Indonesian Borneo. *Science* 303: 1000–1003.  
View Article • PubMed/NCBI • Google Scholar
2. Ancrenaz M, Dabek L, O'Neil S (2007) The cost of exclusion: recognizing a role for local communities in biodiversity conservation. *Plos Biol* 5(11): e289. doi:10.1371/journal.pbio.0050289.  
View Article • PubMed/NCBI • Google Scholar
3. Rijksen HD (1978) A field study on Sumatran orang-utans (*Pongo pygmaeus abelii*, Lesson 1827): Ecology, behavior, and conservation. PhD dissertation, H. Veenman and Zonen, Wageningen, The Netherlands.
4. Rao M, van Schaik CP (1997) The behavioural ecology of Sumatran orang-utans in logged and unlogged forest. *Trop Biod* 4(2): 173–185.  
View Article • PubMed/NCBI • Google Scholar

5. Rijksen HD, Meijaard E (1999) *Our Vanishing Relative: The Status of Wild Orangutans at the Close of the Twentieth Century*. Dordrecht: Kluwer Academic Publishers.. 480 p.
6. Morrogh-Bernard H, Husson SJ, Page SE, Rieley JO (2003) Population status of the Bornean orang-utan (*Pongo pygmaeus*) in the Sabangau peat-swamp forest, Central Kalimantan, Indonesia. *Biol Cons* 110(1): 141–51.  
View Article • PubMed/NCBI • Google Scholar
7. Felton AM, Engstrom LM, Felton A, Knott CD (2003) Orangutan population density, forest structure and fruit availability in hand-logged and unlogged peat swamp forests in West Kalimantan, Indonesia. *Biol Cons* 114: 91–101.  
View Article • PubMed/NCBI • Google Scholar
8. Johnson AE, Knott CD, Pamungkas B, Pasaribu , Marshall AJ (2005) A survey of the orangutan (*Pongo pygmaeus wurmbii*) population in and around Gunung Palung National Park, West Kalimantan, Indonesia based on nest counts. *Biol Cons* 121: 495–507.  
View Article • PubMed/NCBI • Google Scholar
9. Russon AE, Erman A, Dennis R (2001) The population and distribution of orang-utans (*Pongo pygmaeus pygmaeus*) in and around the Danau Sentarum Wildlife Reserve, West Kalimantan, Indonesia. *Biol Cons* 97: 21–28.  
View Article • PubMed/NCBI • Google Scholar
10. Knop E, Ward PI, Wich SA (2004) A comparison of orang-utan density in a logged and unlogged forest on Sumatra. *Biol Cons* 120: 187–192.  
View Article • PubMed/NCBI • Google Scholar
11. Ancrenaz M, Gimenez O, Ambu L, Ancrenaz K, Andau P, et al. (2005) Aerial surveys give new estimates for orang-utans in Sabah, Malaysia. *Plos Biol* 3 (1): 30–37.  
View Article • PubMed/NCBI • Google Scholar
12. Meijaard E, Sheil D (2007) The persistence and conservation of Borneo's mammals in lowland rain forests managed for timber: observations, overviews and opportunities. *Ecol Res* 23: 21–34. *Ecol S Japan DOI* 10.1007/s11284-007-0342-7.  
View Article • PubMed/NCBI • Google Scholar
13. Van Kreveld A, Roerthorst (2009) *Great Apes and Logging*. Zeist, Switzerland: World Wide Fund Report.
14. Husson S, Wich S, Marshall AJ, Dennis RD, Ancrenaz M, et al. (2009) Orangutan distribution, density, abundance and impacts of disturbance. In: Wich S, Utami S, Setia T, van Schaik C, editors. *Geographic variation in behavioral ecology and conservation*. Oxford University Press. pp. 77–96.
15. Wich SA, Meijaard E, Marshall AE, Husson S, Ancrenaz M, et al. (2008) Distribution and conservation status of the orang-utan (*Pongo spp.*) on Borneo and Sumatra: how many remain? *Oryx* 42 (3): 1–11.  
View Article • PubMed/NCBI • Google Scholar
16. Sabah Forestry Department (2008) *Forest management Plan for Ulu Segama Malua Sustainable Forest Project Area*. Sandakan, , Sabah,, Malaysia..
17. Mathewson P, Spehar S, Meijaard E, Nardiyono , Purnomo , Sasmirul A, Sudiyanto , Oman , Sulhudin , Jasary , Jumali , Marshall AJ (2008) Evaluating orangutan census techniques using nest decay rates: implications for population estimates. *Ecol Applications* 18: 208–221.  
View Article • PubMed/NCBI • Google Scholar
18. Kuehl H, Maisel F, Ancrenaz M, Williamson EA (2008) *Best practice guidelines for surveys and monitoring of great ape populations*. Gland, Switzerland: IUCN/SSC Primate Specialist Group.
19. Ancrenaz M, Gimenez O, Goossens B, Sawang A, Lackman-Ancrenaz I (2004) Determination of ape distribution and population size with ground and aerial surveys: a case study with orang-utans in lower Kinabatangan, Sabah, Malaysia. *Anim Cons* 7: 375–385.  
View Article • PubMed/NCBI • Google Scholar
20. Boyko RH, Marshall AJ (2010) Using simulation models to evaluate ape nest survey techniques. *Plos One* in press.
21. Marsh CW (1995) *Danum Valley Conservation Area, Sabah, Malaysia: management plan*. Kota Kinabalu Yayasan Sabah/Innoprise Sdn Bhd.
22. Harrison T, Harrison B (1971) The prehistory of Sabah. *Sabah Soc J* 1–272.  
View Article • PubMed/NCBI • Google Scholar
23. Pryer WB (1883) Notes on North-eastern Borneo and the Sulu islands. *Proc Royal Geog Soc* 5: 90–96.  
View Article • PubMed/NCBI • Google Scholar
24. Marshall AJ, Nardiyono , Engström LM, Pamungkas B, Palapa J, Meijaard E, Stanley SA (2006) The blowgun is mightier than the chainsaw in determining population density of Bornean orangutans (*Pongo pygmaeus morio*) in the forests of East Kalimantan. *Biol Cons* 129: 566–578.  
View Article • PubMed/NCBI • Google Scholar
25. Marshall AJ, Lacy R, Ancrenaz M, Byers O, Husson S, et al. (2009) Orangutan population biology, life history, and conservation: Perspectives from PVA models. In: Wich S, Utami S, Setia T, van Schaik C, editors. *Geographic variation in behavioral ecology and conservation*. Oxford University Press. pp. 311–326.

26. Marshall AJ, Salas LA, Stephens S, Nardiyono, Engström LM, Meijaard E, Stanley S (2007) Use of limestone karst forests by Bornean orangutans (*Pongo pygmaeus morio*) in the Sangkulirang Peninsula, East Kalimantan, Indonesia. *Am J Primatol* 69: 1–8.  
View Article • PubMed/NCBI • Google Scholar
27. Goossens B, Chikhi L, Jalil F, Ancrenaz M, Lackman-Ancrenaz I, et al. (2005) Patterns of genetic diversity and migration in increasingly fragmented and declining orang-utan (*Pongo pygmaeus*) populations from Sabah, Malaysia. *Mol Ecol* 14: 441–456.  
View Article • PubMed/NCBI • Google Scholar
28. Goossens B, Setchell JM, James S, Funk SM, Chikhi L, et al. (2006) Philopatry and reproductive success in Bornean orang-utans (*Pongo pygmaeus*). *Mol Ecol* 15: 2577–2588.  
View Article • PubMed/NCBI • Google Scholar
29. McKinnon JR (1974) The behavior and ecology of wild orangutans (*Pongo pygmaeus*). *Anim Behav* 22: 3–74.  
View Article • PubMed/NCBI • Google Scholar
30. Payne J (1988) Orang-utan conservation in Sabah. Kuala Lumpur: WWF-Malaysia International, Report 3759.
31. Djojosedharmo S, van Schaik CP (1992) Why are orang utans so rare in the highlands?: Altitudinal changes in a Sumatran forest. *Trop Biod* 1: 11–22.  
View Article • PubMed/NCBI • Google Scholar
32. Russon AE, Wich SA, Ancrenaz M, Kanamori T, Knott CD, et al. (2009) Geographic variation in orangutan diets. In: Wich S, Utami S, Setia T, van Schaik C, editors. *Geographic variation in behavioral ecology and conservation*. Oxford University Press. pp. 135–156.
33. Marshall AJ, Ancrenaz M, Brearley FQ, Fredriksson GM, Ghaffar N, et al. (2009b) The effects of forest phenology and floristics on populations of Bornean and Sumatran orangutans. In: Wich S, Utami S, Setia T, van Schaik C, editors. *Geographic variation in behavioral ecology and conservation*. Oxford University Press. pp. 97–118.
34. Singleton I, van Schaik CP (2001) Orangutan home range size and its determinants in a Sumatran swamp forest. *Int J Prim* 22: 877–911.  
View Article • PubMed/NCBI • Google Scholar
35. Johns AD (1988) Effects of “selective” timber extraction on rain forest structure and composition and some consequences for frugivores and folivores. *Biotropica* 20(1): 31–7.  
View Article • PubMed/NCBI • Google Scholar
36. Morrogh-Bernard H, Husson S, Knott CD, Wich SA, van Schaik CP, et al. (2009) Orangutan activity and diet, a comparison between species, populations and habitats. In: Wich S, Utami S, Setia T, van Schaik C, editors. *Geographic variation in behavioral ecology and conservation*. Oxford University Press. pp. 119–134.
37. MacKinnon K, Hatta G, Halim H, Mangalik A (1996) *The ecology of Kalimantan*. Singapore: Periplus Editions.
38. Siegert F, Ruecker G, Hinrichs A, Hoffman AA (2001) Increased damages from fires in logged forests during droughts caused by El Nino. *Nature* 414: 437–440.  
View Article • PubMed/NCBI • Google Scholar
39. Taylor AB (2009) The functional significance of variation in jaw form in orangutans. In: Wich S, Utami S, Setia T, van Schaik C, editors. *Geographic variation in behavioral ecology and conservation*. Oxford University Press. pp. 15–32.
40. van Schaik CP, Marshall AJ, Wich SA (2009) Geographic variation in orangutan behavior and biology. In: Wich S, Utami S, Setia T, van Schaik C, editors. *Geographic variation in behavioral ecology and conservation*. Oxford University Press. pp. 351–361.
41. Knott CD (1998) Changes in orangutan caloric intake, energy balance, and ketones in response to fluctuating fruit availability. *Int J Primatol* 19: 1061–1079.  
View Article • PubMed/NCBI • Google Scholar
42. Meijaard E, Sheil D, Nasi DR, Augeri D, Rosenbaum B, et al. (2005) *Life After Logging: Reconciling Wildlife Conservation and Production Forestry in Indonesian Borneo*. Jakarta, Indonesia: Center for International Forestry Research.. 345 p.
43. Sheil D, Meijaard E (2010) Purity and prejudice deluding ourselves about biodiversity conservation. In press.
44. Wilson K, Meijaard E, Drummond H, Boitani L, Catullo G, et al. (2010) Conserving biodiversity in production landscapes. *Ecol Applic*. In press.  
View Article • PubMed/NCBI • Google Scholar
45. Walsh RPD (1996) Drought frequency changes in Sabah and adjacent parts of northern Borneo since the late nineteenth century and possible implications for tropical rain forest dynamics. *J Trop Ecol* 12(3): 385–407.  
View Article • PubMed/NCBI • Google Scholar
46. Van Schaik CP, Azwar A, Priatna D (1995) Population estimates and habitat preferences of orangutans based on line transects of nests. In: Nadler RD, Galdikas BMF, Sheeran LK, Rosen N, editors. *The Neglected Ape*. New York: Plenum Press. pp. 129–147.

47. Cassey P, McArdle H (1999) An assessment of distance sampling techniques for estimating animal abundance. *Environmetrics* 10: 261–278.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

48. Buckland ST, Anderson DR, Burnham KP, Laake JL (1993) *Distance Sampling: Estimating Abundance of Biological Populations*. London: Chapman and Hall.. 446 p.