

Climate change is projected to reduce carrying capacity and redistribute species richness in North Pacific pelagic marine ecosystems

PHOEBE A. WOODWORTH-JEFCOATS^{1,2}, JEFFREY J. POLOVINA¹ and JEFFREY C. DRAZEN²
¹NOAA Fisheries, Pacific Islands Fisheries Science Center, 1845 Wasp Blvd., Bldg. 176, Honolulu, HI 96818, USA, ²School of Ocean and Earth Science and Technology, University of Hawaii at Manoa, 1000 Pope Road, Marine Sciences Building, Honolulu, HI 96822, USA

Abstract

Climate change is expected to impact all aspects of marine ecosystems, including fisheries. Here, we use output from a suite of 11 earth system models to examine projected changes in two ecosystem-defining variables: temperature and food availability. In particular, we examine projected changes in epipelagic temperature and, as a proxy for food availability, zooplankton density. We find that under RCP8.5, a high business-as-usual greenhouse gas scenario, increasing temperatures may alter the spatial distribution of tuna and billfish species richness across the North Pacific basin. Furthermore, warmer waters and declining zooplankton densities may act together to lower carrying capacity for commercially valuable fish by 2–5% per decade over the 21st century. These changes have the potential to significantly impact the magnitude, composition, and distribution of commercial fish catch across the pelagic North Pacific. Such changes will in turn ultimately impact commercial fisheries' economic value. Fishery managers should anticipate these climate impacts to ensure sustainable fishery yields and livelihoods.

Keywords: carrying capacity, climate change impacts, commercial fisheries, Coupled Model Intercomparison Project 5, North Pacific, pelagic habitat, zooplankton

Received 4 May 2016; revised version received 8 August 2016 and accepted 9 August 2016

Introduction

Pelagic marine habitat is projected to experience a number of impacts from climate change (e.g., Bopp *et al.*, 2013). As earth system models improve with each model generation, confidence in their projections has increased and a community consensus is coalescing around several projected impacts. Of these, two of the most significant impacts to epipelagic habitat are likely to be ocean warming (Bopp *et al.*, 2013) and the expansion of the oligotrophic subtropical gyres (Sarmiento *et al.*, 2004; Steinacher *et al.*, 2010; Polovina *et al.*, 2011; Cabré *et al.*, 2015). Ocean warming is a direct result of ocean heat uptake in response to atmospheric warming driven by increasing greenhouse gas concentrations. Gyre expansion is projected as the result of two physical mechanisms. Ocean heating leading to increased vertical stratification is expected to further reduce nutrient concentrations in the euphotic zone of oligotrophic gyre waters (Xu *et al.*, 2012; Cabré *et al.*, 2015). Additionally, changes in atmospheric circulation may result in a poleward displacement of both the descending branch of the Hadley circulation and midlatitude storm

tracks (Chang *et al.*, 2012; Scheff & Frierson, 2012; Yongyun *et al.*, 2012; Cabré *et al.*, 2015). These changes in atmospheric circulation will in turn alter ocean surface wind stress curl, primarily along the gyres' poleward boundaries, contributing to gyre expansion.

Change in the biophysical marine environment will impact many marine organisms, as well as fisheries and those who rely on fishery services. Catch from pelagic fisheries in the North Pacific is largely comprised of tuna, including skipjack (*Katsuwonus pelamis*), yellowfin (*Thunnus albacares*), and bigeye (*Thunnus obesus*), as well as other species (FAO, 2012). Tunas occupy specific thermal habitats at different life stages and have a high metabolic demand (Lehodey *et al.*, 2011, 2013). Thus, changes to either thermal habitat or ocean productivity are likely to impact tunas, as well as other commercially valuable fish. We examine these climate change impacts through a suite of models included in the fifth phase of the Coupled Model Intercomparison Project (CMIP5; Taylor *et al.*, 2012), focusing on the two habitat parameters that most directly influence ecosystem productivity and capacity: temperature and food availability. We aim to make broad projections of climate change impacts on marine fisheries that can be used by fishery managers when drafting ecosystem-based fisheries management plans. Previous studies

Correspondence: Phoebe A. Woodworth-Jefcoats, tel. +1 (808) 725 5562, fax +1 (808) 725 5474, e-mail: Phoebe.Woodworth-Jefcoats@noaa.gov

have suggested that climate change may have a substantial impact on commercial fish catch, independent of fishing (Cheung *et al.*, 2010; Lehodey *et al.*, 2011, 2013; Bell *et al.*, 2013; Howell *et al.*, 2013; Woodworth-Jefcoats *et al.*, 2015). Therefore, it is essential that these potential impacts be incorporated into management plans so that both fishery resources and livelihoods can be sustained well into the future.

Materials and methods

Earth system models used

We examine 11 earth system models included in CMIP5. Models used are presented in Table 1. Models selected are those with two trophic levels (phyto- and zooplankton) of output available at time of download. All data were downloaded from the CMIP5 data portal (http://cmip-pcmdi.llnl.gov/cmip5/data_portal.html). Spherical interpolation (for curvilinear grids) and nearest coordinate regridding (for rectilinear grids) were used to regrid output to a common $1^\circ \times 1^\circ$ rectilinear grid spanning $0\text{--}66^\circ\text{N}$ and $120^\circ\text{E}\text{--}70^\circ\text{W}$, with the Bering Sea and Sea of Okhotsk excluded. We note that output from two additional models, HadGEM2-CC and HadGEM2-ES (Collins *et al.*, 2011), was available but not used in our analysis due to unrealistic negative plankton densities across much of the central North Pacific.

Data used

Both historical and projected data are examined. All projections are from the representative concentration pathway (RCP) 8.5 scenario, 'a relatively conservative business as usual case... with no explicit climate policy' (Riahi *et al.*, 2011). We focus on two 20-year time periods representing the beginning and end of the 21st century: 1986–2005 and 2081–2100. The beginning of the 21st century is captured by the last 20 years of the historical runs and the end of the 21st century by the last 20 years of the 21st century in the RCP8.5 projection.

Data are vertically integrated across the epipelagic zone, represented as the upper 200 m of the water column. Vertical resolution varies by model, and we integrated across all depths of 200 m or less. We examine potential temperature, phytoplankton carbon density, and zooplankton carbon density output by each model. Data are examined as a model ensemble to address the possible influence of individual model drift (Sen Gupta *et al.*, 2013).

Vertically integrated (upper 200 m) ocean temperatures from the World Ocean Atlas 2013 (WOA13; Locarnini *et al.*, 2013) were used as baseline temperatures when calculating projected ecosystem impacts. The temperature increase projected by each model was then added to the WOA13 data to determine projected ecosystem change.

Pelagic habitat

We examine changes in thermal habitat by comparing probability frequency distributions of pooled 20-year, monthly

epipelagic temperatures. Because so much of the literature is focused on SST, we also present projected changes in SST though these changes are not the focus of our analysis. In both cases, monthly temperatures were used in an effort to fully capture seasonal extremes and distributions are binned in 0.5°C bins. Change in zooplankton densities is similarly compared, although annual densities are used as this is the only temporal scale available for three-dimensional biogeochemical data through the CMIP5 data portal. Distributions are binned in 0.05 g C m^{-2} bins. Twenty-year means from the beginning and end of the 21st century are used to evaluate the absolute change in epipelagic temperature and percent change in both phyto- and zooplankton densities.

Ecosystem impacts

We assess two measures of ecosystem impact: tuna and billfish species richness and carrying capacity. Species richness captures the total number of tuna and billfish species present and carrying capacity the total number of fish the ecosystem can support. Species richness (SR) is a function of epipelagic temperature, following Eqn (1) as determined by Boyce *et al.* (2008),

$$SR = -0.0033T^3 + 0.1156T^2 - 0.4675T \quad (1)$$

where T is epipelagic temperature in $^\circ\text{C}$. Carrying capacity (K) is determined from ecological theory, following Eqn (2),

$$K \propto [R]M^{-3/4}e^{E/kT} \quad (2)$$

where $[R]$ is limiting resource supply, which we take as zooplankton density, M is target fish mass, E is activation energy (0.63 eV; Brown *et al.*, 2004), k is Boltzmann's constant ($8.62 \times 10^{-5}\text{ eV K}^{-1}$; Brown *et al.*, 2004; Jennings *et al.*, 2008), and T is epipelagic temperature in Kelvin (Brown *et al.*, 2004). Given that Eqn (2) is a proportional relationship, we evaluate relative changes in the right-hand side of the equation and refer to these as changes in potential carrying capacity (K_p). Twenty-year means from the beginning and end of the 21st century are used to evaluate changes in SR and K_p . We hold M constant over both periods so the resulting change in K_p is independent of M . To assess whether R or T has a greater influence on K_p , we examine the difference between the absolute percent change in both R and $e^{E/kT}$ following Eqn (3),

$$|\%\Delta R| - |\%\Delta e^{E/kT}| \quad (3)$$

with positive results indicating that changes in zooplankton density have the greatest influence on K_p and negative results indicating that changes in T have the greatest influence on K_p .

Results

Pelagic habitat

Thermal habitat. Across all models, sea surface and epipelagic temperatures are projected to increase (Table 1, Figs 1a and 2a). Additionally, all model scenarios project the emergence of new, warmer temperatures by the end of the 21st century. Emerging SSTs

Table 1 For each model, the SST and epipelagic temperature ranges that decrease in frequency, increase in frequency, and emerge by the end of the 21st century followed by the zooplankton density ranges that increase and decrease in frequency by the end of the century

Model	Sea surface temperature (°C)			Epipelagic temperature (°C)			Zooplankton density (g C m ⁻²)	
	Decreases	Increases	Emerges	Decreases	Increases	Emerges	Increases	Decreases
Canadian Centre for Climate Modelling and Analysis Earth system model* (CanESM2)	20.0–30.0	30.0–38.0	34.0–38.0	14.5–21.5	21.5–32.5	30.5–32.5	0.05–0.20	0.20–0.40
NOAA Geophysical Fluid Dynamics Laboratory Earth System Model Generalized ocean layer dynamics† (GFDL-ESM2G)	20.0–29.5	29.5–35.0	32.5–35.0	13.0–18.5	24.5–32.0	30.0–32.0	0.50–0.90	0.90–1.85
NOAA Geophysical Fluid Dynamics Laboratory Earth System Model Modular Ocean Model 4† (GFDL-ESM2M)	20.0–30.0	30.0–35.0	33.0–35.0	15.0–25.5	25.5–32.5	30.0–32.5	0.40–0.95	0.95–1.75
NASA Goddard Institute for Space Studies ModelE2 Earth System Model with carbon cycle coupled to the HYCOM ocean model‡§ (GISS-E2-H-CC)	22.0–30.0	30.0–34.5	32.0–34.5	17.5–23.0	23.0–32.5	31.0–32.5	0.00–0.10	0.10–1.00
NASA Goddard Institute for Space Studies ModelE2 Earth System Model with carbon cycle coupled to the Russell ocean model‡§ (GISS-E2-R-CC)	20.0–30.5	30.5–34.5	32.5–34.5	16.5–26.5	26.5–33.0	31.5–33.0	0.00–0.15	0.15–0.85
Institut Pierre-Simon Laplace Low Resolution CM5A¶ (IPSL-CM5A-LR)	21.0–30.0	30.0–36.0	32.5–36.0	15.5–26.0	26.0–34.0	30.0–34.0	0.30–0.45	0.60–1.10
Institut Pierre-Simon Laplace Medium resolution CM5A¶ (IPSL-CM5A-MR)	21.5–31.0	31.0–36.5	33.0–36.5	15.5–26.0	26.0–34.0	31.0–34.0	0.30–0.65	0.65–0.95
Institut Pierre-Simon Laplace Low resolution CM5B¶ (IPSL-CM5B-LR)	21.5–30.0	30.0–35.5	32.0–35.5	17.5–24.0	26.5–32.5	31.0–32.5	0.25–0.35	0.35–0.70
Max-Planck-Institute für Meteorologie Earth System Model low resolution** (MPI-ESM-LR)	20.0–30.0	30.0–37.0	34.0–37.0	14.5–22.0	22.0–33.5	31.5–33.5	0.10–0.75	0.75–1.50
Max-Planck-Institute für Meteorologie Earth System Model medium resolution** (MPI-ESM-MR)	20.0–30.0	30.0–36.5	34.0–36.5	16.0–21.0	21.0–35.0	32.5–35.0	0.10–0.70	0.70–1.55
Meteorological Research Institute Earth System Model Version 1†† (MRI-ESM1)	21.0–29.5	21.0–34.5	31.5–34.5	16.5–27.0	27.0–30.5	29.0–30.5	0.00–0.20	0.20–0.40

*Christian *et al.* (2010).†Dunne *et al.* (2013).‡Romanou *et al.* (2014).§Schmidt *et al.* (2014).¶Dufresne *et al.* (2013).**Giorgetta *et al.* (2013).††Yukimoto *et al.* (2011).

(i.e., temperatures not present at the beginning of the 21st century that are present at the end of the 21st century) range from 31.5 to 38 °C and epipelagic

temperatures from 29 to 35 °C. Change in thermal habitat is also captured through the difference between frequency distributions over time. Epipelagic




Fig. 1 Percent frequency distributions of 20-year pooled epipelagic temperatures (a) and zooplankton densities (b) from the beginning (gray) and end (green) of the 21st century are plotted above the difference between the two distributions. Solid lines show multimodel means. Terciles encompassing 33%, 67%, and 100% of the models are shaded progressively lighter. The red line in the lower panel of (a) indicates the temperature range over which at least half the models project the emergence of new thermal habitat.

temperatures that comprise the majority of the North Pacific at the beginning of the 21st century (15.6–23.7 °C on average) decline in frequency and warmer temperatures come to dominate by the end of the century (24.5–32.9 °C on average).

Our results focus on the warmest temperatures in the North Pacific as these temperatures cover the largest area. However, it is important to note that there is a similar distributional shift in the coolest temperatures. Here, too, there is model consensus on a shift toward warmer temperatures, as well as a loss of the coolest temperatures by the end of the 21st century (Fig. 1a). Across all models, disappearing epipelagic temperatures range from -1.5 to 2 °C. Three models (IPSL-CM5B-LR, MPI-ESM-LR, and MPI-ESM-MR) project a loss of the coolest SSTs, ranging from -2.0 to -1.0 °C.

Food available to fish. We take zooplankton density to be a proxy for food available to fish. Across all models, the distribution of zooplankton densities is projected to shift toward lower values (Table 1, Figs 1b and 2b). Densities that comprise the majority of the North Pacific at the beginning of the 21st century (0.50 – 1.10 g C m $^{-2}$ on average) decline in frequency and lower densities come to dominate by the end of the century (0.18 – 0.49 g C m $^{-2}$ on average).

Not only do the models used in our study project zooplankton densities to decline across much of the North Pacific, but they also project these declines to be amplified relative to declines in phytoplankton densities (Fig. 2c, warm colors represent waters where zooplankton declines are projected to be greater than phytoplankton declines). When declining zooplankton densities are examined in relation to projected phytoplankton changes, we find that zooplankton declines exceed phytoplankton declines to a large degree. All models but three (CanESM2, GISS-E2-H-CC, and GISS-E2-R-CC) place such waters across much of the North

Pacific excluding only subpolar waters, and in some cases equatorial and California Current upwelling waters. Projected declines in zooplankton exceed those of phytoplankton by 10–30% on average, with individual model maxima of 25–50% found along the periphery of the North Pacific subtropical gyre (NPSG).

Ecosystem impacts

Changes in predicted tuna and billfish species richness (SR) follow projected changes in epipelagic temperature. Across all models, the area of maximum SR shifts northward and eastward. Species richness declines across much of the central and western subtropics and increases in temperate and subpolar waters, with the magnitude of change increasing with distance toward the western tropical Pacific and temperate latitudes, peaking at approximately four species lost or gained (Fig. 2d). Most models project potential carrying capacity (K_p) for commercially valuable fish to decline by 20–50% across the North Pacific, or by roughly 2–5% per decade over the 21st century (Fig. 2e). As with trophic amplification, the areas projected to see the greatest declines in K_p are found along the periphery of the NPSG. Declining K_p is a result of both increasing epipelagic temperature and declining zooplankton density, with the primary driver varying across the North Pacific. In the western equatorial Pacific and NPSG, declining zooplankton density has a stronger impact on K_p , while in the eastern equatorial Pacific and at temperate latitudes, increasing epipelagic temperature is the stronger driver (Fig. 2f).

Discussion

The CMIP5 projections presented in this study suggest a number of changes to North Pacific pelagic habitat. Broadly, thermal habitat is projected to warm and be




Fig. 2 Multimodel median projected change in epilagic habitat (a and b) and resulting degrees of ecosystem impact (c–f) over the 21st century: projected change in epilagic temperature (a) and zooplankton density (b), degree of trophic amplification (indicated by warm colors) or the difference between projected phytoplankton and zooplankton percent declines (c), projected change in tuna and billfish species richness for waters within the bounds of a positive solution to Eqn (1) (5–30 °C) (d), projected percent change in potential carrying capacity (e), and the difference in the strength of changing zooplankton density (warm colors) vs. changing epilagic temperature (cool colors) as drivers of change in potential carrying capacity (f). In (a–e), stippling indicates areas where at least 80% of the models used project a change of the same sign. In (f), stippling indicates areas where at least 80% of the models used indicate the same dominant driver.

spatially redistributed. Zooplankton densities are projected to decline and to an amplified degree relative to phytoplankton declines. When these projections are examined more finely and in relation to one another, they suggest that commercial fisheries in the central North Pacific may see catch decline by 20–50% and be comprised of three to four fewer tuna and billfish species.

Changing pelagic habitat

While warming epipelagic temperatures might be expected to unfold as a straightforward poleward creep of present-day conditions, we find that this is not the case (Fig. 2a, d). Rather, warmer temperatures appear to emerge from the western equatorial Pacific and expand eastward and northward as moderate temperatures retreat in kind. Over time, this results in a reshaping of pelagic thermal habitat. For example, thermal habitat associated with adult tuna foraging is displaced by thermal habitat more commonly associated with tuna spawning grounds, and spawning habitat is replaced by temperatures that exceed even the warmest temperatures associated with commercially valuable fish (Boyce *et al.*, 2008; Lehodey *et al.*, 2011, 2013).

Evidence suggests that fish and other pelagic organisms will relocate to maintain residence in preferred thermal habitat in both freshwater (Grenouillet & Comte, 2014) and marine (Pinsky *et al.*, 2013; Montero-Serra *et al.*, 2015) environments, and with relocations varying over different life history stages (Walsh *et al.*, 2015). Some fish may simply be able to spend more time in deeper, cooler waters. However, such an adaptation comes at a cost. For example, fish may forage less successfully at the lower light levels found below the epipelagic realm. Organisms that are unable to exploit deeper habitat will be forced to relocate geographically. Such vertical and geographic relocations could ultimately alter predator–prey dynamics.

The emergence of new thermal habitat also raises questions, as it is projected to exceed current maximum temperatures. It remains unknown how or whether pelagic organisms will adapt to these temperatures. Storch *et al.* (2014) suggest there are firm limits on temperatures to which animals can adapt. They find that due to constraints posed by cellular complexity, the highest SST that allowed multicellular Eukaryota to grow was 40 °C, close to temperatures projected to occur over the North Pacific in our study. The unprecedented rate at which climate is changing (Doney *et al.*, 2014) adds further uncertainty to questions surrounding adaptation.

In addition to changes in thermal habitat, we also project a shift toward lower zooplankton densities over

the 21st century. Spatially, the lowest zooplankton densities are associated with the oligotrophic NPSG. Declining densities are manifested as both an expansion of the NPSG and lower densities in NPSG waters (Fig. 2b). While we examine the oligotrophic NPSG from the perspective of zooplankton densities, our results are similar to those from other studies focused on phytoplankton that project the gyre's expansion (Sarmiento *et al.*, 2004; Steinacher *et al.*, 2010; Polovina *et al.*, 2011; Cabré *et al.*, 2015).

Not only do the models used in our study project zooplankton densities to decline across much of the North Pacific, but they also project these declines to be amplified relative to declines in phytoplankton densities. Stock *et al.* (2014) link trophic amplification to declining zooplankton growth efficiency as food resources (net primary production) decline, while Chust *et al.* (2014) link trophic amplification to nonlinear coupling of phytoplankton and zooplankton biomass. It remains unclear whether this amplification in the plankton community will propagate further up through the food web; however, modeling work suggests that it will be amplified by some micronekton (Bell *et al.*, 2013) and possibly throughout the size spectrum (Lefort *et al.*, 2015). If trophic amplification does indeed carry through the food web, an amplification of roughly 20% at each trophic linkage could result in apex predator density (trophic level 4–5) declining by up to 50–60% by the end of the century, or by 5–6% per decade.

Ecosystem impacts of changing pelagic habitat

The projected impacts of climate change in the North Pacific extend beyond the immediate changes to temperature and food availability. Increasing epipelagic temperature is projected to lead to a redistribution of tuna and billfish SR (Fig. 2d). There is strong model agreement of a decline of up to 3–4 species across much of the subtropics with an increase of similar magnitude projected for temperate latitudes. These projected changes in SR largely, and not surprisingly given Eqn (1), mirror the changing footprint of thermal habitat in the North Pacific. Based solely on thermal tolerance, much of the subtropical North Pacific is projected to become less hospitable to adult commercially valuable tuna and billfish. While a decline of only a few species may not seem very substantial, the longline fisheries in these waters target only a small number of species, primarily bigeye tuna and swordfish (*Xiphias gladius*), and also catch several commercially valuable, nontarget species such as skipjack tuna, yellowfin tuna, shortbill spearfish (*Tetrapturus anguistorostris*), and striped marlin (*Tetrapturus audax*). Thus, even a small

decline in SR could significantly impact catch composition, magnitude, and value. Likewise, at the northern limits of the fishery, the small increase in species diversity could potentially benefit fishermen. Whether this potential benefit would be offset by the increased expense of traveling further from port to fish is unknown. Fishermen may also shift their homeport based on target catch relocation, as some in the Hawaii-based longline fishery have already done.

Increasing epipelagic temperatures combined with largely declining zooplankton densities are projected to act together to lower North Pacific K_p over the 21st century. We find strong model agreement that K_p is projected to decline by roughly 20–50% across the North Pacific (Fig. 2e). Despite our measure of K_p being a simple relationship based on ecological theory, this projection is in line with previous studies that have projected similar declines in exploitable high-trophic-level biomass as the result of climate change (Lefort *et al.*, 2015; Woodworth-Jefcoats *et al.*, 2015). We also find that declines in K_p exceed those of zooplankton densities, further suggesting that trophic amplification in the plankton community may propagate up through the food web. Additionally, K_p is projected to decline even in regions where plankton densities are projected to increase (Fig. 2b, e). This suggests that potential increases in biomass at the base of the food web would not be enough to compensate for the metabolic costs of increasing temperatures. Further examination of the impact of temperature vs. zooplankton on K_p shows that the dominant driver of change varies spatially (Fig. 2f). In subtropical regions where zooplankton declines are projected to be greatest, these declines seem to have the greater impact on K_p . In the eastern North Pacific and at temperate latitudes, waters seeing the greatest increase in epipelagic temperature, temperature increases drive K_p declines.

Potential carrying capacity is projected to decline most in and around the central North Pacific. This has the potential to particularly impact longline fisheries operating in this area. Potential fisheries' yields could decline by up to 50% over a time when the Food and Agriculture Organization of the United Nations projects that food resources will need to increase by roughly 70% to meet the demands of a growing human population (UN, 2011). Such an increase in demand could further strain the ecosystem, as the heavy removal of large fish has the potential to drive down exploitable biomass independent of any bottom-up impacts (Blanchard *et al.*, 2005; Ward & Myers, 2005; Polovina & Woodworth-Jefcoats, 2013).

The areas of greatest trophic amplification and declining K_p occur around the boundaries of the NPSG (Fig. 2c, e). To the north of the NPSG lies the North

Pacific transition zone, a narrow area used as a migration and foraging corridor by a number of pelagic species (Polovina *et al.*, 2001; Hazen *et al.*, 2013). To the south of the NPSG are spawning grounds for a number of tropical tuna species (Lehodey *et al.*, 2011, 2013). Thus, the areas likely to see the greatest declines in food availability are areas crucial to specific life history stages of pelagic species. Such a mismatch in resource demand and supply could amplify climate impacts on species exploiting these regions. Furthermore, given that organisms from around the North Pacific target these areas, changes here have the potential to impact the entire basin. These maxima of declining phyto- and zooplankton densities are not flanked by corresponding areas of increasing densities, suggesting that productive regions around the NPSG are not simply relocating. Or, if productive regions are relocating, they are still experiencing overall declines in phytoplankton densities. The importance of these regions bordering the NPSG, along with their relatively small size, makes them ideal areas for monitoring climate change as it unfolds. Survey (Howell *et al.*, 2015; Polovina *et al.*, 2015) and tagging (Block *et al.*, 2011) efforts already in place in these regions may provide insight into how organisms across the food web are responding to climate change.

One question we are unable to address in this study is how regions bordering the NPSG may be impacted by changes in phenology. The transition zone in particular moves meridionally with the seasons. The phenology of both the seasonal migration of the transition zone (Hazen *et al.*, 2013) and its associated productivity (Polovina *et al.*, 2011) may change as a result of climate change. Thus, organisms targeting the region at specific times of the year may have to migrate farther or to different locations. Both finer temporal resolution projections and tagging data may help address such phenology questions.

Caveats

Our study focuses on the two primary influences on ecosystem capacity: temperature and food availability. These are far from the only influences, however. Other variables such as oxygen concentration, pH, and exploitation can influence pelagic carrying capacity. Given that changes in many of these variables are projected to have negative impacts in the North Pacific (Koslow *et al.*, 2011; Bopp *et al.*, 2013), they are likely to exacerbate the impacts of warming temperatures and declining food availability.

We also assume that physical climate influences will be the primary determinants of ecosystem capacity. However, species and trophic interactions are also

influential. In some cases, these interactions can have a larger impact than physical climate drivers (Grenouillet & Comte, 2014; Ockendon *et al.*, 2014). Additionally, changes in temperature and food availability can alter foraging range and create new competition (Bond & Lavers, 2014). Such changes in predator–prey interactions could have large impacts on commercial fisheries and could potentially be examined through species-based ecosystem modeling approaches and network theory.

In this study, we examine only the epipelagic realm, although many commercially valuable fish also inhabit mesopelagic depths (Howell *et al.*, 2010; Abecassis *et al.*, 2012). Future impact studies could examine a broader vertical habitat range. For example, Lefort *et al.* (2015) suggest that fishes able to migrate between epipelagic and mesopelagic depths may fare better in the face of climate change than fishes restricted to either realm. Finally, we examine only one climate change scenario. By examining RCP8.5, we hopefully project the upper limits of potential climate change impacts. Future work could examine more optimistic RCPs, potentially providing motivation to take mitigating actions by presenting goals for limited impacts.

Commercial fishery impacts of changing pelagic habitat

Through examining a suite of CMIP5 earth system models, we find that climate change may significantly alter North Pacific epipelagic habitat over the 21st century. Warming thermal habitat and declining zooplankton densities are projected to lower potential carrying capacity, and in turn fishery yield, by approximately 2–5% per decade. Additionally, based on changing thermal habitat alone, species richness across much of the subtropics is projected to decline by up to four tuna and billfish species by the end of the century. Together, these changes have the potential to significantly impact commercial fish catch in the North Pacific. Fishery managers can use these projections to place current yields and management actions in a broader climate-based context. For example, early warning thresholds for changing catch composition or yield could be based on projected climate impacts. Such strategic management plans would ensure that the ecosystem is not further stressed by unsustainable removals.

Acknowledgements

We thank Axel Timmermann, Evan Howell, and an anonymous reviewer for insights that greatly improved this manuscript. We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modeling groups (listed in Table 1 of this article) for producing and making available their

model output. For CMIP in the US Department of Energy's Program for Climate Model Diagnosis and Intercomparison provide coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth System Science Portals. We also acknowledge use of the Ferret program for analysis in this study. Ferret is a product of NOAA's Pacific Marine Environmental Laboratory. (Information is available at <http://ferret.pmel.noaa.gov/Ferret/>) This is SOEST contribution number 9741.

References

- Abecassis M, Dewar H, Hawn D, Polovina J (2012) Modeling swordfish daytime vertical habitat in the North Pacific Ocean from pop-up archival tags. *Marine Ecology Progress Series*, **452**, 219–236.
- Bell JD, Ganachaud A, Gehrke PC *et al.* (2013) Mixed responses of tropical Pacific fisheries and aquaculture to climate change. *Nature Climate Change*, **3**, 591–599.
- Blanchard JL, Dulvy NK, Jennings S, Ellis JR, Pinnegar JK, Tid A, Kell LT (2005) Do climate and fishing influence size-based indicators of Celtic Sea fish community structure? *ICES Journal of Marine Science*, **62**, 405–411.
- Block BA, Jonsen ID, Jorgensen SJ *et al.* (2011) Tracking apex marine predator movements in a dynamic ocean. *Nature*, **475**, 86–90.
- Bond AL, Lavers JL (2014) Climate change alters the trophic niche of a declining, apex marine predator. *Global Change Biology*, **20**, 2100–2107.
- Bopp L, Resplandy L, Orr JC *et al.* (2013) Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models. *Biogeosciences*, **10**, 6225–6245.
- Boyce DG, Tittensor DP, Worm B (2008) Effects of temperature on global patterns of tuna and billfish richness. *Marine Ecology Progress Series*, **355**, 267–297.
- Brown JH, Gillooly JF, Allen AP, Savage VM, West GB (2004) Toward a metabolic theory of ecology. *Ecology*, **85**, 1771–1789.
- Cabr  A, Marinov I, Leung S (2015) Consistent global responses of marine ecosystems to future climate change across the IPCC AR5 earth system models. *Climate Dynamics*, **45**, 1253–1280.
- Chang EKM, Guo Y, Xia X (2012) CMIP5 multimodel ensemble projection of storm track change under global warming. *Journal of Geophysical Research*, **117**, D23118.
- Cheung WWL, Lam VWY, Sarmiento JL, Kearney K, Watson R, Zeller D, Pauly D (2010) Large-scale redistribution of maximum fisheries catch potential in the global ocean under climate change. *Global Change Biology*, **16**, 24–35.
- Christian JR, Arora VK, Boer CJ *et al.* (2010) The global carbon cycle in the Canadian Earth system model (CanESM1): preindustrial control simulation. *Journal of Geophysical Research*, **115**, G03014.
- Chust W, Allen JL, Bopp L *et al.* (2014) Biomass changes and trophic amplification of plankton in a warmer ocean. *Global Change Biology*, **20**, 2124–2139.
- Collins WJ, Bellouin N, Doutriaux-Boucher M *et al.* (2011) Development and evaluation of an Earth-system model – HadGEM2. *Geoscientific Model Development*, **4**, 1051–1075.
- Doney SC, Bopp M, Long MC (2014) Historical and future trends in ocean climate and biogeochemistry. *Oceanography*, **27**, 108–119.
- Dufresne J-L, Foujols M-A, Denvil S *et al.* (2013) Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5. *Climate Dynamics*, **40**, 2123–2165.
- Dunne JP, John JG, Shevliakova E *et al.* (2013) GFDL's ESM2 global coupled climate-carbon Earth System Models. Part II: carbon system formulation and baseline simulation characteristics. *Journal of Climate*, **26**, 2247–2267.
- Food and Agriculture Organization of the United Nations (FAO) (2012) *FAO Yearbook, Fishery and Aquaculture Statistics, Capture Production*. FAO, Rome, Italy.
- Giorgetta MA, Jungclauss J, Reick CH *et al.* (2013) Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5. *Journal of Advances in Modeling Earth Systems*, **5**, 572–597.
- Grenouillet G, Comte L (2014) Illuminating geographical patterns in species' range shifts. *Global Change Biology*, **20**, 3080–3091.
- Hazen EL, Jorgensen S, Rykaczewski RR *et al.* (2013) Predicted habitat shifts of Pacific top predators in a changing climate. *Nature Climate Change*, **3**, 234–238.
- Howell EA, Hawn DR, Polovina JJ (2010) Spatiotemporal variability in bigeye tuna (*Thunnus obesus*) dive behavior in the central North Pacific Ocean. *Progress in Oceanography*, **86**, 81–93.
- Howell EA, Wabnitz CCC, Dunne JP, Polovina JP (2013) Climate-induced primary productivity change and fishing impacts on the Central North Pacific ecosystem and Hawaii-based pelagic longline fishery. *Climatic Change*, **119**, 79–93.

- Howell EA, Bograd SJ, Hoover AL, Seki MP, Polovina JJ (2015) Variation in phytoplankton composition between two North Pacific frontal zones along 158°W during winter–spring 2008–2011. *Progress in Oceanography*. doi: 10.1016/j.pocean.2015.06.003.
- Jennings S, Mélin F, Blanchard JL, Forster RM, Dulvy NK, Wilson RW (2008) Global-scale predictions of community and ecosystem properties from simple ecological theory. *Proceedings of the Royal Society B*, **275**, 1375–1383.
- Koslow JA, Goericke R, Lara-Lopez A, Watson W (2011) Impact of declining intermediate-water oxygen on deepwater fishes in the California Current. *Marine Ecology Progress Series*, **436**, 207–218.
- Lefort S, Aumont O, Bopp L, Arsouze T, Gehlen M, Maury O (2015) Spatial and body-size dependent response of marine pelagic communities to projected global climate change. *Global Change Biology*, **21**, 154–164.
- Lehodey P, Hampton J, Brill RW *et al.* (2011) Vulnerability of oceanic fisheries in the tropical Pacific to climate change. In: *Vulnerability of Tropical Pacific Fisheries and Aquaculture to Climate Change* (eds Bell JD, Johnson JE, Hobday AJ), pp. 433–492. Secretariat of the Pacific Community, Noumea, New Caledonia.
- Lehodey P, Semoma I, Calmettes B, Hampton J, Nicol S (2013) Modeling the impact of climate change on Pacific skipjack tuna population and fisheries. *Climatic Change*, **119**, 95–109.
- Locarnini RA, Mishonov AV, Antonov JI *et al.* (2013) *World Ocean Atlas 2013, Volume 1: Temperature*. Levitus S, Ed., Mishonov A, Technical Ed., NOAA Atlas NESDIS 73. National Oceanographic Data Center, Silver Spring, MD, USA.
- Montero-Serra I, Edwards M, Genner MJ (2015) Warming shelf seas drive the subtropicalization of European pelagic fish communities. *Global Change Biology*, **21**, 144–153.
- Ockendon N, Baker DJ, Carr JA *et al.* (2014) Mechanisms underpinning climatic impacts on natural populations: altered species interactions are more important than direct effects. *Global Change Biology*, **20**, 2221–2229.
- Pinsky ML, Worm B, Fogarty MJ, Sarmiento JL, Levin SA (2013) Marine taxa track local climate velocities. *Science*, **341**, 1239–1242.
- Polovina JJ, Woodworth-Jefcoats PA (2013) Fishery-induced changes in the subtropical Pacific pelagic ecosystem size structure: observations and theory. *PLoS One*, **8**, e62341.
- Polovina JJ, Howell E, Kobayashi DR, Seki MP (2001) The transition zone chlorophyll front, a dynamic global feature defining migration and forage habitat for marine resources. *Progress in Oceanography*, **49**, 469–483.
- Polovina JJ, Dunne JP, Woodworth PA, Howell EA (2011) Projected expansion of the subtropical biome and contraction of the temperature and equatorial upwelling biomes in the North Pacific under global warming. *ICES Journal of Marine Science*, **68**, 986–995.
- Polovina JJ, Howell E, Kobayashi DR, Seki MP (2015) The transition zone chlorophyll front updated: advances from a decade of research. *Progress in Oceanography*. doi: 10.1016/j.pocean.20105.01.006.
- Riahi K, Rao S, Krey V *et al.* (2011) RCP 8.5 – a scenario of comparatively high greenhouse gas emissions. *Climatic Change*, **109**, 33–57.
- Romanou A, Romanski J, Gregg WW (2014) Natural ocean carbon cycle sensitivity to the parameterizations of the recycling in a climate model. *Biogeosciences*, **11**, 1137–1154.
- Sarmiento JL, Slater R, Barber R *et al.* (2004) Response of ocean ecosystems to climate warming. *Global Biogeochemical Cycles*, **18**, GB3003.
- Scheff J, Frierson DMW (2012) Robust future precipitation declines in CMIP5 largely reflect the poleward expansion of model subtropical dry zones. *Geophysical Research Letters*, **39**, L18704.
- Schmidt GA, Kelley M, Nazarenko L *et al.* (2014) Configuration and assessment of GISS ModelE2 contributions to the CMIP5 archive. *Journal of Advances in Modeling Earth Systems*, **6**, 141–184.
- Sen Gupta A, Jourdain NC, Brown JN, Monselesan D (2013) Climate drift in the CMIP5 models. *Journal of Climate*, **26**, 8597–8615.
- Steinacher M, Joos F, Frölicher TL *et al.* (2010) Projected 21st century decrease in marine productivity: a multi-model analysis. *Biogeosciences*, **7**, 979–1005.
- Stock CA, Dunne JP, John JG (2014) Drivers of trophic amplification of ocean productivity trends in a changing climate. *Biogeosciences*, **11**, 7125–7135.
- Storch D, Menzel L, Frickenhaus S, Pörtner H-O (2014) Climate sensitivity across marine domains of life: limits to evolutionary adaptation shape species interactions. *Global Change Biology*, **20**, 3059–3067.
- Taylor KE, Stouffer RJ, Meehl GA (2012) An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, **93**, 485–498.
- United Nations (UN) (2011) Seven billion and growing: the role of population policy in achieving sustainability. Technical Paper No. 2011/13. 36 pp.
- Walsh HJ, Richardson DE, Marancik KE, Hare JA (2015) Long-term changes in the distributions of larval and adult fish in the Northeastern U.S. shelf ecosystem. *PLoS One*, **10**, e0137382.
- Ward P, Myers RA (2005) Shifts in open-ocean fish communities coinciding with the commencement of commercial fishing. *Ecology*, **86**, 835–847.
- Woodworth-Jefcoats PA, Polovina JJ, Howell EA, Blanchard JL (2015) Two takes on the ecosystem impacts of climate change and fishing: comparing a size-based and a species-based ecosystem model in the central North Pacific. *Progress in Oceanography*, **138**, 533–545.
- Xu L, Xie S-P, Lui Q (2012) Mode water ventilation and subtropical countercurrent over the North Pacific in CMIP5 simulations and future projections. *Journal of Geophysical Research*, **117**, C12009.
- Yongyun H, Lijun T, Jiping L (2012) Poleward expansion of the Hadley circulation in CMIP5 simulations. *Advances in Atmospheric Sciences*, **30**, 790–795.
- Yukimoto S, Yoshimura H, Hosaka M *et al.* (2011) Meteorological Research Institute–Earth System Model Version 1 (MRI-ESM1) Model Description. Technical Report of the Meteorological Research Institute No. 64.